



ELSEVIER

Linear Algebra and its Applications 294 (1999) 9–33

**LINEAR ALGEBRA
AND ITS
APPLICATIONS**

www.elsevier.com/locate/laa

Finiteness of a set of non-collinear vectors generated by a family of linear operators

R. Mubarakzjanov ^{*,1}

Department of Theoretical Cybernetics, Kazan University, 420008 Kazan, Russian Federation

Received 3 December 1997; accepted 29 January 1999

Submitted by S. Friedland

Abstract

Let \mathbb{R}^n be a real n -dimensional space, let $\{A(x) \mid x \in X\}$ be a family of $m = |X|$ linear operators in \mathbb{R}^n , and let K_r be a sharp polyhedral cone formed by a set of r vectors, $K_r \subset \mathbb{R}^n$. Let K_r be invariant under $\{A(x) \mid x \in X\}$, i.e. $K_r A(x) = K_r$, for $x \in X$. We study a maximum set of non-collinear vectors derived from a vector $\mathbf{h} \in K_r$ by the family $\{A(x) \mid x \in X\}$ in this paper. It is shown that there is a function $f(n, m, r)$ such that this set of non-collinear vectors is finite iff the cardinality of this set is not greater than $f(n, m, r)$. This result can be used for solving the following problem: when does a channel simulated by a probabilistic automaton have a finite set of states? © 1999 Elsevier Science Inc. All rights reserved.

Keywords: Linear operator; Eigenvectors; Polyhedral cone; Probabilistic automaton

1. Introduction

A channel, simulated by a finite probabilistic automaton [1], can be described by an initial vector and a family of matrices. In [1, p. 66, Theorem 2.4], an algorithm is given to decide whether the set of channel states is finite. But it

^{*} Corresponding author. E-mail: rustam.mubarakzjanov@ksu.ru

¹ This paper has been written during scholarship in the Rostock University, Germany (Stipend of German Academic Exchange Service (DAAD)). Research partially supported by the Russian Fund for Basic Research 96-01-01692.